



## CULTIVACE MAGNESIUM

### GUARANTEED ANALYSIS

Magnesium (Mg) ..... 5.0%

**DERIVED FROM:** Magnesium Acetate

**CAUTION: KEEP OUT OF THE REACH OF CHILDREN, HARMFUL IF SWALLOWED**

*This product is intended to prevent or correct a magnesium deficiency.*

10.0 lb/gal @ 68°F

### COMPATIBILITY

This product is compatible with most pesticides, however, it is not possible to test all pesticide mixes. A jar test is recommended before use.

### GENERAL USE RECOMMENDATIONS

**DILUTION:** Field and row crops: 3 to 20 gallons by air, 10 to 40 gallons by ground. Tree crops: concentrate 50 to 100 gallons, dilute 100 to 500 gallons.

### NET CONTENTS 2.5 GALLONS (9.46 L)

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>



### PRECAUTIONARY STATEMENTS

**Danger :** Causes severe skin burns and eye damage.

**Prevention :** Wear protective gloves. Wear eye or face protection. Wear protective clothing. Wash hands thoroughly after handling.

**Response :** IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. Immediately call a POISON CENTER or physician. IF SWALLOWED: Immediately call a POISON CENTER or physician. Rinse mouth. Do NOT induce vomiting. IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water or shower. Wash contaminated clothing before reuse. Immediately call a POISON CENTER or physician. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or physician.

**Storage :** Store locked up.

**Disposal :** Dispose of contents and container in accordance with all local, regional, national and international regulations.

Guaranteed by KWS Distributing LLC.  
17887 SE Grand Island Rd.  
Dayton, Oregon 97114

**SUGGESTED USES:** (rates per acre) Crops such as, but not limited to:

**Alfalfa, hay and other forage crops :** 1 - 4 quarts - After early spring growth and after cutting.

**Apples, cherries, peaches, almonds, walnuts, pears, stone, pome fruits, citrus and nuts :** 1 - 4 quarts per acre at 2 to 4 week intervals.

**Barley, oats, wheat, rice, and other small grains :** 1 - 4 quarts - Apply at tillering, early boot and flag leaf emergence.

**Beans, peas and other legumes :** 1 - 3 quarts - Apply at early bud, repeat after bloom.

**Cabbage, broccoli, carrots, onions, tomatoes and other vegetables :** 1 - 4 quarts - Apply when growth is 4 - 8" high. Repeat every 2 weeks as needed.

**Corn, field and sweet :** 1 - 4 quarts - Apply when 10 - 20" high, repeat in 2 - 4 weeks as needed.

**Sugar Cane:** 1 - 4 pints per acre.

**Grapes :** 1 - 4 quarts - Apply dormant, pre-bloom and post-bloom as needed.

**Hops :** 1 - 4 quarts

**Lettuce, celery, spinach and other leafy vegetables :** 1 - 4 quarts - Apply after transplanting or thinning, repeat as needed.

**Hazelnuts, pistachios, other deciduous trees :** 1 - 4 quarts - Apply at bud break Repeat every two trees to three weeks.

**Potatoes :** 1 - 4 quarts - Apply at tuber initiation, repeat in 2 - 4 week intervals.

**Raspberries, Blueberries, strawberries, Blackberries :** 1 - 3 quarts - Apply pre bloom repeat every 2-4 weeks.

**Grasses grown for seed :** 1-4 quarts. Apply at tillering, early boot and flag leaf emergence.

**NOTICE** Manufacturer makes no warranties, expressed or implied. The buyer assumes all risks associated with the use of this product. In the event of damage resulting from a breach of warranty, the buyer agrees to accept a refund of the purchase price as full discharge of the manufacturers liability.

Rev. 070517

**CULTIVACEGROWTH.COM**  
A KWS DISTRIBUTING COMPANY