



Research Driven,
Proven Results®

GRIGG® CARBOPLEX®

6-4-4

GRIGG Carboplex contains three sources of nitrogen, phosphate, potassium, amino acid complexed micronutrients and sea plant extract (*Ascophyllum nodosum*). It helps boost turf strength, vigor and hardiness – making it a great option for winter turfgrass preparation.

Key Advantages

- Contains soluble nutrients for efficient uptake and use
- Nitrogen promotes consistent turfgrass shoot growth
- Phosphorus plays a role in plant metabolic processes that transfer energy throughout the plant
- Potassium regulates primary physiological processes that impact turf response to stress and supports cellular processes that impact photosynthesis, water regulation, respiration and protein production
- Iron plays a key role in chlorophyll production, which improves turf color

Application and Use

Foliar Applications: Apply as needed every 7-14 days.

Tees, Greens: 6-12 fl oz per 1,000 sq ft or 2-4 gal per acre [20-40 L per hectare].

Fairways, Sports Field: 9-12 fl oz per 1,000 sq ft or 3-4 gal per acre [30-40 L per hectare].

Soil Applications: Apply as needed every 7-21 days.

Tees, Greens: 9-12 fl oz per 1,000 sq ft or 3-4 gal per acre [30-40 L per hectare].

Fairways, Sports Fields: 12-15 fl oz per 1,000 sq ft or 4-5 gal per acre [40-50 L per hectare].

**For a distributor near you contact:
800 300 6559 or www.grigg.co**

GRIGG is part of Brandt Consolidated, Inc.
2935 South Koke Mill Road
Springfield, IL 62711
www.brandt.co

Guaranteed Analysis

Total Nitrogen (N)	6.00%
0.70% Ammoniacal nitrogen	
1.10% Nitrate nitrogen	
4.20% Urea nitrogen	
Available Phosphate (P ₂ O ₅)	4.00%
Soluble Potash (K ₂ O)	4.00%
Iron (Fe)	0.20%
0.20% Water soluble iron	
Manganese (Mn)	0.05%
0.05% Water soluble manganese	
Zinc (Zn)	0.05%
0.05% Water soluble zinc	

Derived from monoammonium phosphate, potassium nitrate, urea, iron amino acid complex, manganese amino acid complex, zinc amino acid complex, potassium hydroxide and kelp (*Ascophyllum nodosum*).

Make frequent applications at lower rates, or apply higher rates for greater plant demand. Optimum rate of application will vary depending on treatment interval, soil properties (such as pH, organic matter content, texture), weather conditions, time of year, plant species and its nutrient requirements. For best results, follow soil/tissue test recommendation.

This product is not intended for use on food crop sites.

